

Reply to comment by Hillis *et al.* (2013)

N. J. White,¹ L. M. Mackay,² S. M. Jones,³ J. P. B. Lovell¹ and M. W. Davis¹

¹*Bullard Laboratories, Madingley Rise, Madingley Road, Cambridge, CB3 0EZ, UK. E-mail: njw10@cam.ac.uk*

²*BP Exploration Ltd., Chertsey Road, Sunbury-on-Thames, Middlesex, TW16 7LN, UK*

³*School of Geography, Earth & Environmental Sciences, University of Birmingham, B15 2TT, UK*

Accepted 2013 April 3. Received 2013 February 20

We agree that there is now convincing evidence which favours the idea that a region encompassing the British Isles is dynamically supported by convective circulation beneath the lithospheric plate. For example, the coincidence of a long wavelength (>800 km) free-air gravity anomaly with anomalously slow velocities beneath the lithospheric plate suggests that regional elevation of Northern Britain is maintained by a tongue of hot asthenosphere protruding away from Iceland (Jones *et al.* 2002a; Arrowsmith *et al.* 2005). These observations are corroborated by the crustal thickness measurements of Davis *et al.* (2012). The Icelandic plume initiated 62 Ma and there is excellent evidence in the Irmingier and Icelandic basins that this plume has waxed and waned with different periodicities (Jones *et al.* 2002b; Poore *et al.* 2009). This transient activity has manifest itself in geochemical and earthquake records along the Reykjanes Ridge (Poore *et al.* 2011; Parnell-Turner *et al.* 2013), in the Neogene record of deep-water overflow across the Greenland-Scotland Ridge (Poore *et al.* 2006), and along the fringing continental margins where spectacular buried ephemeral landscapes have been encountered (Shaw-Champion *et al.* 2008; Rudge *et al.* 2008; Hartley *et al.* 2011). Around the British Isles, there is considerable onshore and offshore evidence favouring a history of transient and/or permanent vertical motions which may reflect Palaeogene and Neogene plume activity (White & Lovell 1997; Jones *et al.* 2001; Jones *et al.* 2002a; Al-Kindi *et al.* 2003; Jones & White 2003; Mackay *et al.* 2005). This regional pattern of epeirogeny is locally complicated by faulting and folding, which is often difficult to date with precision.

Hillis *et al.* (2013) have raised two issues which challenge the significance of convectively driven epeirogeny. First, they state that our ‘map of Cenozoic denudation is . . . highly inaccurate’. Secondly, they argue that horizontal shortening alone can account for the pattern of regional denudation. It is important to understand how the denudation map shown in Davis *et al.* (2012) was constructed. First, >3000 sonic velocity, seismic velocity, subsidence and fission track estimates of denudation were assembled. We omitted denudation estimates obtained by inverting vitrinite reflectance data because they have excessive scatter (Rowley & White 1998). Even so, the database is noisy on short wavelengths. Some of this scatter may be geologically meaningful since regional uplift and denudation are undoubtedly moderated by shorter wavelength faulting and folding. Nonetheless, all four methods for estimating denudation rarely agree and individually suffer from significant uncertainties which are unlikely to be smaller than ± 500 m. These difficulties encouraged us to calculate mean values of denudation in $0.5^\circ \times 1^\circ$ (i.e. 50×50 km) bins. These mean values were then contoured using a minimum curvature interpolating surface with a tension factor of 0.15. Hillis *et al.* (2013) point out that Fig. 10 indicates 0–0.5 km in the Weald basin and on the East Midlands Shelf where they

have published spot measurements based on sonic velocity analyses which are >2 km (Hillis *et al.* 2008). Their assertion is incorrect. In the Weald basin, we show a northward embayment which has 0.5–1 km of denudation. On the East Midlands Shelf, several bands have 0.5–1 km of denudation and higher values of 1–1.5 km occur at Flamborough Head.

It is instructive to scrutinize denudation estimates from the Weald basin and from the East Midlands Shelf in more detail. In the Weald basin, Hillis *et al.* (2008) used sonic velocity analyses to argue for denudation values of >2.2 km along the axis of the Weald anticline. They have not quoted uncertainties. In an earlier study, Butler & Pullam (1990) used a combination of stratigraphic reconstructions and interval velocity measurements to propose that up to 1.5 km of Late Cretaceous and Cenozoic strata have locally been removed from the crest of the Weald anticlinorium. Their Fig. 5(b) shows interval velocities which are anomalously fast but subject to considerable scatter. Jones (1999) published a revised stratigraphic reconstruction which implies that up to 1.1–1.4 m are missing from the Central Weald. These different estimates have uncertainties of ± 500 m. Inverse modelling of subsidence data from the Weald basin suggests that denudation is 0.1–1 km. It is even less clear when denudation occurred and to what extent regional (i.e. longer wavelength) epeirogeny played a role (Blundell 2002). The consensus is that a series of discrete episodes occurred in Palaeogene, Neogene and Plio-Pleistocene times when extensive peneplains developed. It is important to stress that Fig. 10 only shows average denudation values, which were binned into 50×50 km blocks of 0.5–1 km, which accord with the range of published estimates.

On the East Midlands Shelf, denudation estimates have been made by analysing fission track and sonic velocity data (e.g. Green 1989; Hillis 1993). Originally, Lewis *et al.* (1992) argued, based on Green’s (1986, 1989) fission track analyses, that up to 3 km of denudation occurred across northern England in Early Cenozoic times. Holliday (1993) suggested that these estimates were excessive because they could not be reconciled with stratigraphic constraints. He argued that 1.2–1.75 km had been eroded from northern England. Green *et al.* (2001) reanalysed the fission track database and concluded that there were two episodes of cooling, at 65–55 Myr and at 25–0 Myr. They indicated that it was not possible to determine the relative proportions of Palaeogene and Neogene denudation. Fission track measurements from four wells on the East Midlands Shelf reveal a west–east gradient in denudation. In our database, we have included denudation estimates from Bray *et al.* (1992), Green *et al.* (2001) and Green (2002) which were corrected, where necessary, using the new assumptions of Green *et al.* (2001). From west to east, these estimates are: (i) Rufford-1 = 1.45 km with a range of 0.91–1.65 km; (ii) Grove-3 = 1.09 km; (iii) Biscathorpe =

460 m and (iv) Cleethorpes-1 = 320 m. Hillis (1993) used sonic velocity analyses to show that the Cleethorpes-1 well was denuded by ~ 1.4 km which agreed with Green's (1989) original fission track estimate of 1.3–1.7. Across the East Midlands Shelf, fission track and sonic velocity estimates of denudation are generally higher than those obtained by inverting subsidence data which suggest only 0–1 km (Rowley & White 1998). Thus averaged fission track estimates show that the East Midlands Shelf was denuded by 0.83 ± 0.5 km. At Flamborough Head itself, Hillis *et al.* (2008) proposed that denudation is ~ 2 km. Stewart & Bailey (1996) compiled different estimates, which show that ~ 1 km of denudation occurred to the south of Flamborough Head. Fig. 10 shows that denudation at Flamborough Head is 1–1.5 km. We agree with Hillis (1993) who stated that 'widely recognized inversion axes are only relatively local culminations of more regional Cenozoic erosion. The base level of apparent erosion in these areas, on which erosion of inversion axes are superimposed, is approximately 1 km'.

Our map of regional denudation, which is based upon a large database of sonic velocity, interval velocity, subsidence and fission track measurements, is consistent with published estimates. Note that long wavelength denudation is greatest in Northern Britain. Nonetheless, Hillis *et al.* (2013) hold great store by minor discrepancies at the crest of the Weald Anticline and at Flamborough Head. We believe that the reason for these discrepancies stems from their reliance upon sonic velocity analyses which form the centrepiece of Hillis *et al.* (2008). This methodology has two important drawbacks. First, an undisturbed compaction trend as a function of depth must be identified against which deviations in sonic velocity measurements are gauged. Secondly, sonic velocity measurements are inherently noisy. Both of these drawbacks give rise to substantial uncertainties in estimates of overburden removal (± 1 km). Since Hillis *et al.* (2008) did not quote uncertainties, it is impossible to gauge the reliability of their estimates or to compare them in a meaningful way with independent estimates of denudation. Mackay & White (2006) have described and applied an integrative methodology for estimating denudation from sonic velocity measurements which is more robust than the standard approach used by Hillis *et al.* (2008). Their methodology yields stable results which do not rely upon estimating an undisturbed compaction trend.

The second issue raised by Hillis *et al.* (2013) concerns the relationship between horizontal shortening and regional denudation. They strongly object to our assertion that 'while there is copious field evidence for phases of horizontal shortening during Cenozoic times, the magnitude of shortening is one order of magnitude smaller than required'. This issue was first addressed quantitatively by Brodie & White (1995) who readily acknowledged that horizontal shortening occurred throughout a region encompassing the British Isles. It is straightforward to calculate the amount of regional shortening that is required to produce a given amount of denudation since the NW European Shelf has a very small elastic thickness (Tiley *et al.* 2003). If a reference column of continental lithosphere is shortened uniformly by a factor, f , then the amount of regional uplift is given by

$$U = \frac{a [(\rho_m - \rho_c) \frac{t_c}{a} (1 - \frac{\alpha T_1}{2} \frac{t_c}{a}) - \frac{\alpha T_1}{2} \rho_m]}{\rho_m (1 - \alpha T_1)} (f - 1), \quad (1)$$

where values of the different symbols are given in Table 1. For a region encompassing the British Isles, it is reasonable to assume that the thicknesses of crust and lithosphere, which yields zero elevation with respect to the mid-oceanic ridge, are $t_c = 30$ km and

Table 1. Symbols and values of parameters used in text are taken from Brodie & White (1995).

Symbol	Parameter	Value
U	Regional uplift	km
D	Regional denudation	km
a	Lithospheric thickness	125 km
t_c	Crustal thickness	30 km
ρ_w	Density of water	1 Mg m ⁻³
ρ_s	Density of sediment	2.4 Mg m ⁻³
ρ_c	Density of crust (at 0°C)	2.8 Mg m ⁻³
ρ_m	Density of lithospheric mantle (at 0°C)	3.33 Mg m ⁻³
ρ_a	Density of asthenosphere	3.2 Mg m ⁻³
α	Thermal expansion coefficient	$3.28 \times 10^{-5} \text{ }^\circ\text{C}^{-1}$
T_1	Asthenospheric temperature	1333°C

$a = 125$ km, respectively. Substitution of the values given in Table 1 yields

$$U = 2.5(f - 1). \quad (2)$$

Since erosion isostatically amplifies regional uplift, the maximum amount of regional denudation is given by

$$D = \left(\frac{\rho_a}{\rho_a - \rho_s} \right) U. \quad (3)$$

If the average density of the sedimentary pile, $\rho_s = 2.4$ Mg m⁻³ then a horizontal shortening, f , of 1.15–1.3 is required to achieve a regional denudation of 1.5–3 km. Given that the British Isles is ~ 1000 km long, we expect to see evidence for 130–230 km of horizontal shortening over this distance. There are at most ~ 7 documented belts of Cenozoic shortening in a region encompassing the British Isles (i.e. Wessex/Weald, South Celtic Sea, St. George's Channel, Sole Pit, East Irish Sea, Moray Firth, Wyville-Thompson/Lui/Westray). Each of these belts is required to have shortened by 19–33 km. Across three of the southernmost structures, Blundell (2002) reported a north–south shortening of ~ 4 km over a distance of 200 km. If we assume that small-scale deformation could double this amount of shortening to ~ 8 km, then $f = 1.02$ – 1.04 which is more than ten times too small. Hillis *et al.* (2013) statement that '15 per cent of shortening is sufficient to generate ~ 1.5 km of denudation' is not correct since their percentage calculation entirely depends upon a specified horizontal distance.

Excellent examples of folding and thrust faulting occur from the England Channel to the Faroe-Shetland basin. Nevertheless, conservative estimates of shortening are at least one order of magnitude (i.e. ten times) less than required. Attempts have been made to argue that substantial amounts of shortening are hidden within apparently undeformed sedimentary rocks. For example, Holford *et al.* (2009) suggested that 'cryptic inversion' can occur whereby large thicknesses of apparently undeformed shale take up shortening by a bulk pure shear mechanism. This suggestion is at odds with three important observations. First, widespread development of pressure solution cleavage and bulk pure shear within these shales is not observed either in outcrop or well data. A useful comparison is southeast France where an ancient continental margin has also been denuded by ~ 2 km. In this case, denudation is achieved by large-scale tight folding, by growth of arrowhead/harpoon structures, and by pervasive cleavage development. Secondly, in regions of active shortening > 90 per cent of deformation is taken up on discrete thrust faults which slip in earthquakes with substantial moment release. Thirdly, and most importantly, if regional uplift and denudation is mainly generated by horizontal shortening, the crust should be thickened by a factor of 1.15–1.3 and crustal thicknesses

of 34.5–39 km are expected. There is evidence that ~35 km thick crust exists beneath the Midland Valley of Scotland and beneath North Wales. However, there is no evidence for thickened crust beneath either the Weald basin or beneath the East Midland Shelf (Davis *et al.* 2013). These structures are rootless and confined to the upper crust (Blundell 2002).

In summary, we maintain that there is a significant shortening deficit which suggests that regional epeirogenic uplift has played a significant, but not necessarily exclusive, role in sculpting the outcrop pattern of the British Isles. Beneath Northern Britain, denudation is greater and circumstantial evidence exists which links episodic phases of uplift to waxing and waning of the Icelandic plume (Jones *et al.* 2001; Jones & White 2003). In Southern Britain, we agree with Blundell's (2002) suggestion that regional epeirogeny has been moderated by short wavelength folding and faulting.

REFERENCES

- Arrowsmith, S.J., Kendall, M., White, N.J., VanDecar, J.C. & Booth, D.C., 2005. Seismic imaging of a hot upwelling beneath the British Isles, *Geology*, **33**, 345–348.
- Al Kindi, S., White, N., Sinha, M. & England, R., 2003. Crustal trace of a hot convective sheet, *Geology*, **31**, 207–210.
- Blundell, D.J., 2002. Cenozoic inversion and uplift of Southern Britain, in *Exhumation of the North Atlantic Margin: Timing, Mechanisms and Implications for Petroleum Exploration*, Vol. 196, pp. 85–101, eds Doré, A.G., Cartwright, J.A., Stoker, M.S., Turner, J.P. & White, N., Geological Society of London, Special Publication.
- Bray, R.J., Green, P.F. & Duddy, I.R., 1992. Thermal history reconstruction using apatite fission track analysis and vitrinite reflectance: a case study from the UK East Midlands and Southern North Sea, in *Exploration Britain: Geological Insights for the Next Decade*, Vol. 67, pp. 3–25, ed. Hardman, R.F.P., Geological Society, Special Publication.
- Brodie, J. & White, N., 1994. Sedimentary basin inversion due to igneous underplating: northwest European continental shelf, *Geology*, **22**, 147–150.
- Brodie, J. & White, N., 1995. The causes of sedimentary basin inversion, *J. Geol. Soc. London Spec. Publ.*, **99**, 179–188.
- Butler, M. & Pullan, C.P., 1990. Tertiary structures and hydrocarbon entrapment in the Weald basin of southern England, in *Tectonic Events Responsible for Britain's Oil and Gas Reserves*, Vol. 55, pp. 371–391, eds Hardman, R.F.P. & Brooks, J., Geological Society, Special Publication.
- Davis, M.W., White, N.J., Priestley, K.F., Baptie, B.J. & Tilmann, F.J., 2012. Crustal structure of the British Isles and its epeirogenic consequences, *Geophys. J. Int.*, **190**, 705–725.
- Green, P.F., 1986. On the thermo-tectonic evolution of Northern England: evidence from fission track analysis, *Geol. Mag.*, **123**, 493–506.
- Green, P.F., 1989. Thermal and tectonic history of the East Midlands shelf (onshore UK) and surrounding regions assessed by apatite fission track analysis, *J. geol. Soc. Lond.*, **146**, 755–773.
- Green, P.F., 2002. Early Tertiary paleo-thermal effects in Northern England: reconciling results from apatite fission track analysis with geological evidence, *Tectonophysics*, **349**, 131–144.
- Green, P.F., Thomson, K. & Hudson, J.D., 2001. Recognition of tectonic events in undeformed regions: contrasting results from the Midland Platform and East Midlands Shelf, Central England, *J. geol. Soc. Lond.*, **158**, 59–73.
- Hartley, R.A., Roberts, G.G., White, N. & Richardson, C., 2011. Transient convective uplift of an ancient buried landscape, *Nat. Geosci.*, **4**, 562–565.
- Hillis, R.R., 1993. Tertiary erosion magnitudes in the East Midlands Shelf, onshore UK, *J. geol. Soc. Lond.*, **150**, 1047–1050.
- Hillis, R.R. *et al.*, 2008. Cenozoic exhumation of the southern British Isles, *Geology*, **36**, 371–374.
- Hillis, R.R. *et al.*, 2013. Comment on 'Crustal structure of the British Isles and its epeirogenic consequences', *Geophys. J. Int.*, in press.
- Holford, S.P., Turner, J.P., Green, P.F. & Hillis, R.R., 2009. Signature of cryptic sedimentary basin inversion revealed by shale compaction data in the Irish Sea, western British Isles, *Tectonics*, **28**, doi:10.1029/2008TC002359.
- Holliday, D.W., 1993. Mesozoic cover over northern England: interpretation of apatite fission track data, *J. geol. Soc. Lond.*, **150**, 657–660.
- Jones, D.K.C., 1999. On uplift and denudation of the Weald, in *Uplift, Erosion and Stability: Perspectives on Long-Term Landscape Development*, Vol. 162, pp. 25–43, eds Smith, B.J., Whalley, W.B. & Warke, P.A., Geological Society, Special Publication.
- Jones, S., White, N. & Lovell, B., 2001. Cenozoic and cretaceous transient uplift in the Porcupine basin and its relationship to a mantle plume, *Geol. Soc. London Spec. Pub.*, **188**, 345–360.
- Jones, S., White, N., Clarke, B., Rowley, E. & Gallagher, K., 2002a. Present and past influence of the Iceland plume on sedimentation. *Geol. Soc. London Spec. Pub.*, **196**, 13–25.
- Jones, S., White, N. & MacLennan, J., 2002b. V-shaped ridges around Iceland: implications for spatial and temporal patterns of mantle convection. *G-cubed*, **3**, doi:10.1029/2002GC000361.
- Jones, S.M. & White, N.J., 2003. Shape and size of the starting Iceland Plume swell, *Earth planet. Sci. Lett.*, **216**, 271–282.
- Lewis, C.L.E., Green, P.F., Carter, A. & Hurford, A.J., 1992. Elevated K/T palaeotemperatures throughout northwest England: three kilometres of Tertiary erosion? *Earth planet. Sci. Lett.*, **112**, 131–145.
- Mackay, L.M. & White, N.J., 2006. Accurate estimates of the spatial pattern of denudation by inversion of stacking velocity data: an example from the British Isles. *Geochem. Geophys. Geosyst.*, **7**, doi:10.1029/2005GC001192.
- Mackay, L.M., Turner, J., Jones, S.M. & White, N.J., 2005. Cenozoic vertical motions in the Moray Firth Basin associated with initiation of the Iceland plume, *Tectonics*, **24**, TC5004, doi:10.1029/2004TC001683.
- Parnell-Turner, R.E., White, N.J., MacLennan, J., Henstock, T.J., Murton, B.J. & Jones, S.M., 2013. Crustal manifestations of a hot transient pulse beneath the Mid-Atlantic Ridge, *Earth planet. Sci. Lett.*, **363**, 109–120.
- Poore, H.R., Samworth, R., White, N.J., Jones, S.M. & McCave, I.N., 2006. Neogene overflow of northern component water at the Greenland-Scotland Ridge, *Geochem. Geophys. Geosyst.*, **7**, Q06010, doi:10.1029/2005GC001085.
- Poore, H.R., White, N. & Jones, S., 2009. A Neogene chronology of Iceland plume activity from V-shaped ridges, *Earth planet. Sci. Lett.*, **283**, 1–13.
- Poore, H., White, N. & MacLennan, J., 2011. Oceanic circulation and mantle melting controlled by radial flow of hot pulses within Icelandic plume, *Nat. Geosci.*, **4**, 558–561.
- Rowley, E. & White, N., 1998. Inverse modelling of extension and denudation in the East Irish Sea and surrounding areas, *Earth planet. Sci. Lett.*, **161**, 57–71.
- Rudge, J.F., Shaw-Champion, M.E., White, N., McKenzie, D. & Lovell, B., 2008. A plume model of transient diachronous uplift at the Earth's surface, *Earth planet. Sci. Lett.*, **267**, 146–160.
- Shaw-Champion, M.E., White, N.J., Jones, S.M. & Lovell, J.P.B., 2008. Quantifying transient mantle convective uplift: an example from the Faroe-Shetland basin, *Tectonics*, **27**, doi:10.1029/2007TC002106.
- Stewart, S.A. & Bailey, H.W., 1996. The Flamborough Tertiary outlier, UK southern North Sea, *J. geol. Soc. Lond.*, **153**, 163–173.
- Tiley, R., McKenzie, D. & White, N., 2003. The elastic thickness of the British Isles, *J. geol. Soc. Lond.*, **160**, 1–4.
- White, N. & Lovell, J.P.B., 1997. Measuring the pulse of a plume with the sedimentary record, *Nature*, **387**, 888–891.